

The functional equation of the zeta function of a global field

All the references are made to

[W] André Weil, **Basic Number Theory**, Springer, 1967.

Recall that a *global field* is a finite degree extension of $\mathbb{Q}$ or of $\mathbb{F}_p(T)$. Let k be a global field. The completion k_v of k at a finite place v has a largest compact subring; this subring is open and local, and its residual field has finite cardinality q_v . The product of the $(1 - q_v^{-s})^{-1}$ converges for $\Re s > 1$ to a holomorphic function which can be continued to a meromorphic function ζ_k on the whole complex plane (see [W], Equation (6) p. 128 and Definition 8 p. 129). Put

$$Z_k(s) := \left(\pi^{-s/2} \Gamma(s/2) \right)^{r_1} \left((2\pi)^{1-s} \Gamma(s) \right)^{r_2} \zeta_k(s),$$

where r_1 and r_2 denote respectively the numbers of real and imaginary places of k . For each place v of k define the Haar measure β_v on k_v as follows. If k_v is nonarchimedian, let β_v be the Haar measure giving the volume 1 to the largest compact subring; if $k_v = \mathbb{R}$, put $\beta_v := dx$; if $k_v = \mathbb{C}$, set $\beta_v := |dz \wedge d\bar{z}|$. The product of the β_v defines a Haar measure β on the locally compact k -algebra $\mathbb{A}$ of adeles of k . Recall that k is discrete and cocompact in $\mathbb{A}$; in particular the β -covolume $\beta(\mathbb{A}/k)$ of k in $\mathbb{A}$ is a positive real number.

Theorem. *We have*

$$Z_k(1-s) = \beta(\mathbb{A}/k)^{2s-1} Z_k(s)$$

Observations. If k is a number field, then $\beta(\mathbb{A}/k)$ is the square root of the absolute value of the discriminant (see Proposition 7 p. 91 of [W]). If k is of positive characteristic, if q is the cardinality of its largest finite subfield (which exists by Theorem 8 p. 77 of [W]), and if g is its genus, then we have $\beta(\mathbb{A}/k) = q^{g-1}$ (see Corollary 1 p. 100 of [W]).

Proof of the Theorem. Let $\Phi : \mathbb{A} \rightarrow \mathbb{C}$ be the function defined on page 127 of [W], and let $\mathbb{A}^\times$ be the idele group of k . For any continuous morphism ω from $\mathbb{A}^\times$ to $\mathbb{C}^\times$ put

$$Z(\omega, \Phi) := \int_{\mathbb{A}^\times} \Phi(x) \omega(x) d\mu(x),$$

where μ is the measure defined on page 120 of [W]. For any $s \in \mathbb{C}$ and any $x \in \mathbb{A}^\times$ let $\omega_s(x)$ be the adelic norm of x raised to the power s . Since there is a positive constant c_k such that

$$Z_k(s) = c_k Z(\omega_s, \Phi)$$

by Equation (6) page 128 of [W], it suffices to check

$$Z(\omega_{1-s}, \Phi) = \beta(\mathbb{A}/k)^{2s-1} Z(\omega_s, \Phi).$$

Identify $\mathbb{A}$ to its topological dual via a nontrivial character of $\mathbb{A}$ trivial on k as explained in Theorem 3 p. 66 of [W], and equip $\mathbb{A}$ with its selfdual measure, so that the Fourier transform of a continuous integrable function on $\mathbb{A}$ is a well defined function on $\mathbb{A}$. Letting Φ' be the Fourier transform of Φ , we claim

$$Z(\omega_{1-s}, \Phi) = Z(\omega_s, \Phi') = \beta(\mathbb{A}/k)^{2s-1} Z(\omega_s, \Phi).$$

Indeed the first equality follows from Theorem 2 p. 121 of [W], and the second one from the above Observations, Proposition 6 p. 113 of [W], and Equation (7) page 129 of [W]. QED

Pierre-Yves Gaillard, Université Nancy 1, 02.28.05